

BENLYSTA (BELIMUMAB) ORDERS

PATIENT INFORMATION

Name: _____ DOB: _____ Height: _____ Weight: _____
 Address: _____ Phone: _____ Allergies: _____
 City, State, Zip: _____ Email: _____

REFERRAL STATUS

☐ New Referral ☐ Dose or Frequency Change ☐ Order Renewal
 Is this the first dose? ☐ Yes ☐ No, date of last infusion: _____ Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

DIAGNOSIS AND ICD-10 CODE

☐ Autoantibody-Positive, Systemic Lupus Erythematosus (SLE) ICD 10 Code: _____
☐ Other: _____ ICD 10 Code: _____

REQUIRED DOCUMENTATION

☐ This signed order form by the provider ☐ H&P and Clinical/Progress notes supporting primary diagnosis
☐ Patient demographics AND insurance information ☐ Labs and Tests supporting primary diagnosis
☐ Pregnancy Test (if applicable) ☐ ANA (anti-nuclear Ab) and/or anti-dsDNA Test Results

List Tried & Failed Therapies, including duration of treatment:

1)
2)

MEDICATION ORDERS

Initial dosing ☐ Benlysta 10 mg/kg IV over 1 hour at week 0, 2, 4 then every 4 weeks thereafter**
☐ Benlysta _____mg IV over 1 hour at Week 0, 2, 4 then every 4 weeks thereafter

Maintenance dosing ☐ Benlysta 10mg/kg IV over 1 hour every 4 weeks**
☐ Benlysta _____ mg IV over 1 hour every 4 weeks

Refills: ☐ X 6 months ☐ X 1 Year ☐ Other: _____

**Dose may be rounded to nearest vial size within +/- 10%. TO PROHIBIT dose rounding check here ☐

☐ RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline
☐ RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration
☐ RN to flush and lock VAD/CVAD per company protocol:
 Other: _____

PREMEDICATION ORDERS

☐ Acetaminophen 650mg PO prior to infusion ☐ Other: _____
☐ Diphenhydramine 25mg PO prior to infusion ☐ Other: _____

EMERGENCY MEDICATIONS

☐ Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO
 Diphenhydramine 25mg-50mg slow IV push over 2-5 mins
 Acetaminophen 325mg-650mg PO
 Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.3mg IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive
 Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (AIC/AIS only)

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO
 Diphenhydramine 25mg slow IV push over 2-5 mins
 Acetaminophen 325mg PO
 Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

PRESCRIBER INFORMATION

Prescriber Name: _____ NPI Number: _____
 Office Phone: _____ Office Fax: _____
 Prescriber Signature: _____ Date: _____