

PATIENT INFORMATION

Name: _____ DOB: _____ Height: _____ Weight: _____
 Address: _____ Phone: _____ Allergies: _____
 City, State, Zip: _____ Email: _____

REFERRAL STATUS
☐ New Referral

☐ Dose or Frequency Change

☐ Order Renewal

 Is this the first dose? ☐ Yes ☐ No, date of last infusion: _____

 Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

DIAGNOSIS AND ICD 10 CODE
☐ X - linked hypophosphatemia

ICD-10 Code: _____

☐ Tumor-induced osteomalacia

ICD-10 Code: _____

☐ Other disorders of phosphorus metabolism

ICD-10 Code: _____

REQUIRED DOCUMENTATION/TESTING
☐ This signed order form by the provider

☐ Documentation that patient has stopped phos meds and Vit D

☐ Patient demographics AND insurance information

☐ Fasting serum phosphorus concentration should be below the reference range for age prior to initiation of treatment

☐ Clinical/Progress notes supporting primary diagnosis

List Tried & Failed Therapies, including duration of treatment:

1)

2)

MEDICATION ORDERS

Indication		Dosing
XLH	Pediatric (>6 months) <10kg	☐ Crysvita 1 mg/kg SubQ every 2 weeks (rounded to nearest 1mg)
	Pediatric (>6 months) >10kg	☐ Crysvita 0.8 mg/kg SubQ every 2 weeks (rounded to nearest 10mg, Max 90mg)
	Adult	☐ Crysvita 1 mg/kg SubQ every 4 weeks (rounded to nearest 10mg, Max 90mg)
TIO	Pediatric (>2 years)	☐ Crysvita 0.4 mg/kg SubQ every 2 weeks (rounded to nearest 10mg)
	Adult	☐ Crysvita 0.5 mg/kg SubQ every 4 weeks (rounded to nearest 10mg)
	Pediatric or Adult	☐ Crysvita _____ mg/kg SubQ every 2 weeks (rounded to nearest 10mg, Max 180mg)
Other		☐ Crysvita _____ mg SubQ every _____ weeks _____

 Refills*: ☐ X 6 months ☐ X 1 Year ☐ _____ doses ☐ Other: _____

*(if not indicated, order will expire 1 year from date signed)

☐ RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline

☐ RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration

☐ RN to flush and lock VAD/CVAD per company protocol

Other: _____

EMERGENCY MEDICATIONS
☐ Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO

Diphenhydramine 25mg-50mg slow IV push over 2-5 mins

Acetaminophen 325mg-650mg PO

Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated

Epinephrine 0.3mg IM/SQ, may repeat x1

Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (AIC/AIS only)

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO

Diphenhydramine 25mg slow IV push over 2-5 mins

Acetaminophen 325mg PO

Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated

Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1

Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

PRESCRIBER INFORMATION

Prescriber Name: _____ NPI Number: _____

Office Phone: _____ Office Fax: _____

Prescriber Signature: _____ Date: _____