

## PATIENT INFORMATION

Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Height: \_\_\_\_\_ Weight: \_\_\_\_\_  
 Address: \_\_\_\_\_ Phone: \_\_\_\_\_ Allergies: \_\_\_\_\_  
 City, State, Zip: \_\_\_\_\_ Email: \_\_\_\_\_

## REFERRAL STATUS

☐ New Referral

☐ Dose or Frequency Change

☐ Order Renewal

Is this the first dose? ☐ Yes ☐ No, date of last infusion: \_\_\_\_\_

Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

## DIAGNOSIS AND ICD-10 CODE

☐ Severe Eosinophilic Asthma

ICD-10 Code: \_\_\_\_\_

☐ Other: \_\_\_\_\_

ICD-10 Code: \_\_\_\_\_

Does your patient have blood eosinophil counts  $\geq$  150 cells/ $\mu$ L within past 12 months? ☐ Yes ☐ No

## REQUIRED DOCUMENTATION

☐ This signed order form by the provider

☐ H&P and Clinical/Progress notes supporting primary diagnosis

☐ Patient demographics AND insurance information

☐ Labs and Tests supporting primary diagnosis, including blood eosinophil counts

☐ Pulmonary Function Tests

List Tried & Failed Therapies, including duration of treatment:

- 1)
- 2)
- 3)

## MEDICATION ORDERS

Initial Dosing

☐ Fasenra \_\_\_\_\_mg SubQ every 4 weeks for three doses then every 8 weeks thereafter

Maintenance Dosing

☐ Fasenra \_\_\_\_\_mg SubQ every 8 weeks

Refills\*: ☐ X 6 months ☐ X 1 Year ☐ Other: \_\_\_\_\_

\*(if not indicated, order will expire 1 year from date signed)

☐ RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline

☐ RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration

☐ RN to flush and lock VAD/CVAD per company protocol

Other: \_\_\_\_\_

## PREMEDICATION ORDERS

☐ Acetaminophen 650mg PO prior to infusion

☐ Other: \_\_\_\_\_

☐ Diphenhydramine 25mg PO prior to infusion

☐ Other: \_\_\_\_\_

## EMERGENCY MEDICATIONS

☐ Administer the following medications as needed for infusion-related reactions per company protocol:

**Adults (weight >40kg):**

Diphenhydramine 25mg-50mg PO

Diphenhydramine 25mg-50mg slow IV push over 2-5 mins

Acetaminophen 325mg-650mg PO

Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated

Epinephrine 0.3mg IM/SQ, may repeat x1

Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (AIC/AIS only)

**Pediatrics (weight <40kg): (may adjust with weight changes)**

Diphenhydramine 25mg PO

Diphenhydramine 25mg slow IV push over 2-5 mins

Acetaminophen 325mg PO

Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated

Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1

Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

## PRESCRIBER INFORMATION

Prescriber Name: \_\_\_\_\_ NPI Number: \_\_\_\_\_

Office Phone: \_\_\_\_\_ Office Fax: \_\_\_\_\_

Prescriber Signature: \_\_\_\_\_ Date: \_\_\_\_\_