

INFLIXIMAB ORDERS

PATIENT INFORMATION

Name: _____	DOB: _____ Height: _____ Weight: _____
Address: _____	Phone: _____ Allergies: _____
City, State, Zip: _____	Email: _____

REFERRAL STATUS

☐ New Referral
 ☐ Dose or Frequency Change
 ☐ Order Renewal

Is this the first dose? ☐ Yes ☐ No, date of last infusion: _____
 Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

DIAGNOSIS AND ICD 10 CODE

☐ Moderate to Severe Ulcerative Colitis ☐ Moderate to Severe Crohn's Disease ☐ Rheumatoid Arthritis ☐ Ankylosing Spondylitis ☐ Psoriatic Arthritis ☐ Plaque Psoriasis ☐ Diagnosis: _____	ICD-10 Code: _____ ICD-10 Code: _____ ICD-10 Code: _____ ICD-10 Code: _____ ICD-10 Code: _____ ICD-10 Code: _____ ICD-10 Code: _____
---	--

REQUIRED DOCUMENTATION

☐ This signed order form by the provider ☐ Patient demographics AND insurance information	☐ H&P and Clinical/Progress notes supporting primary diagnosis ☐ Labs and Tests supporting primary diagnosis
--	---

MEDICATION ORDERS

Initial Dosing Maintenance Dosing Alternative Dosing	☐ Infliximab _____mg/kg IV at week 0, 2, 6 then every 8 weeks thereafter ☐ Infliximab _____mg/kg IV every 8 weeks ☐ Infliximab _____ IV every _____ weeks ☐ Round to the nearest 100mg vial
--	--

Refills: ☐ X 6 months ☐ X 1 Year Other: _____
*(if not indicated, order will expire 1 year from date signed)

Preferred Brand: ☐ No Preference

☐ Avsola ☐ Inflectra ☐ Remicade ☐ Renflexis

☐ RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline
☐ RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration
☐ RN to flush and lock VAD/CVAD per company protocol
 Other: _____

PREMEDICATION ORDERS

☐ Acetaminophen 650mg PO prior to infusion ☐ Diphenhydramine 25mg PO prior to infusion	☐ Other: _____ ☐ Other: _____
---	--

EMERGENCY MEDICATIONS

☐ Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO
 Diphenhydramine 25mg-50mg slow IV push over 2-5 mins
 Acetaminophen 325mg-650mg PO
 Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.3mg IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (AIC/AIS only)

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO
 Diphenhydramine 25mg slow IV push over 2-5 mins
 Acetaminophen 325mg PO
 Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

PRESCRIBER INFORMATION

Prescriber Name: _____	NPI Number: _____
Office Phone: _____	Office Fax: _____
Prescriber Signature: _____	Date: _____