

PATIENT INFORMATION

Name: _____ DOB: _____ Height: _____ Weight: _____
Address: _____ Phone: _____ Allergies: _____
City, State, Zip: _____ Email: _____

REFERRAL STATUS

☐ New Referral ☐ Dose or Frequency Change ☐ Order Renewal

Is this the first dose? ☐ Yes ☐ No, date of last infusion: _____ Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

DIAGNOSIS AND ICD-10 CODE

☐ Diagnosis: _____ ICD-10 Code: _____

REQUIRED DOCUMENTATION

- ☐ This signed order form by the provider ☐ H&P and Clinical/Progress notes supporting primary diagnosis
☐ Patient demographics AND insurance information ☐ Labs and Tests supporting primary diagnosis
☐ Pregnancy test, as applicable

List Tried & Failed Therapies, including duration of treatment:

- 1)
2)
3)

MEDICATION ORDERS

☐ Keytruda 200 mg IV over 30 minutes once every 3 weeks ☐ Keytruda 400 mg IV over 30 minutes once every 6 weeks

Refills*: ☐ X 6 months ☐ X 1 Year ☐ Other: _____

*(if not indicated, order will expire 1 year from date signed)

- ☐ RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline
☐ RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration
☐ RN to flush and lock VAD/CVAD per company protocol
☐ Other: _____

PREMEDICATION ORDERS

☐ Acetaminophen 650mg PO prior to infusion ☐ Other: _____
☐ Diphenhydramine 25mg PO prior to infusion ☐ Other: _____

EMERGENCY MEDICATIONS

☐ Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO
Diphenhydramine 25mg-50mg slow IV push over 2-5 mins
Acetaminophen 325mg-650mg PO
Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated
Epinephrine 0.3mg IM/SQ, may repeat x1
Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (AIC/AIS only)

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO
Diphenhydramine 25mg slow IV push over 2-5 mins
Acetaminophen 325mg PO
Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated
Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1
Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

PRESCRIBER INFORMATION

Prescriber Name: _____ NPI Number: _____
Office Phone: _____ Office Fax: _____
Prescriber Signature: _____ Date: _____