

PATIENT INFORMATION

Name: _____ DOB: _____ Height: _____ Weight: _____
 Address: _____ Phone: _____ Allergies: _____
 City, State, Zip: _____ Email: _____

REFERRAL STATUS

☐ New Referral ☐ Dose or Frequency Change ☐ Order Renewal

Is this the first dose? ☐ Yes ☐ No, date of last infusion: _____ Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

DIAGNOSIS AND ICD-10 CODE

☐ Myasthenia gravis (gMG)	ICD-10 Code: _____
☐ Paroxysmal Nocturnal Hemoglobinuria (PNH)	ICD-10 Code: _____
☐ Neuromyelitis Optica Spectrum Disorder (NMOSD)	ICD-10 Code: _____
☐ Atypical Hemolytic Uremic Syndrome (aHUS)	ICD-10 Code: _____
☐ Diagnosis _____	ICD-10 Code: _____

REQUIRED DOCUMENTATION

☐ This signed order form by the provider	☐ Antibody results (if applicable): Anti-AQP4 for NMOSD, AChR for gMG
☐ Patient demographics AND insurance information	☐ Documentation of meningococcal vaccines
☐ H&P and Clinical/Progress notes supporting primary diagnosis	

Is your patient enrolled in the Ultomiris-REMS program? ☐ Yes ☐ No (must be enrolled to start therapy)
 Is the ordering PROVIDER enrolled in the Ultomiris and Soliris REMS? ☐ Yes ☐ No (must be enrolled to start therapy)

List Tried & Failed Therapies, including duration of treatment:

1)
2)

MEDICATION ORDERS

Loading Dose	☐ Ultomiris _____ mg IV as a single dose
Maintenance Dose	☐ Ultomiris _____ mg IV every _____ weeks, starting 2 weeks after the loading dose

Infusion rate determined by patient weight in accordance with manufacturer guidelines.

Refills*: ☐ None ☐ X 6 months ☐ X 1 Year ☐ Other: _____

☐ RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline
☐ RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration
☐ RN to flush and lock VAD/CVAD per company protocol
 Other: _____

PREMEDICATION ORDERS

☐ Acetaminophen 650mg PO prior to infusion	☐ Other: _____
☐ Diphenhydramine 25mg PO prior to infusion	☐ Other: _____

EMERGENCY MEDICATIONS

☐ Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO
 Diphenhydramine 25mg-50mg slow IV push over 2-5 mins
 Acetaminophen 325mg-650mg PO
 Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.3mg IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive
 Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (A/C/AIS only)

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO
 Diphenhydramine 25mg slow IV push over 2-5 mins
 Acetaminophen 325mg PO
 Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated
 Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1
 Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

PRESCRIBER INFORMATION

Prescriber Name: _____ NPI Number: _____
 Office Phone: _____ Office Fax: _____
 Prescriber Signature: _____ Date: _____