

PATIENT INFORMATION

Name: _____ DOB: _____ Height: _____ Weight: _____
Address: _____ Phone: _____ Allergies: _____
City, State, Zip: _____ Email: _____

REFERRAL STATUS

☐ New Referral

☐ Dose or Frequency Change

☐ Order Renewal

Is this the first dose? ☐ Yes ☐ No, date of last infusion: _____

Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

DIAGNOSIS AND ICD-10 CODE

☐ Myasthenia Gravis without (acute) exacerbation

ICD-10 Code: _____

☐ Myasthenia Gravis with (acute) exacerbation

ICD-10 Code: _____

☐ Chronic Inflammatory demyelinating polyneuropathy (CIDP)

ICD-10 Code: _____

☐ Other: _____

ICD-10 Code: _____

REQUIRED DOCUMENTATION

☐ This signed order form by the provider

☐ Clinical/Progress notes supporting primary diagnosis

☐ Patient demographics AND insurance information

☐ Labs and Tests supporting primary diagnosis

☐ MGFA Classification and MG ADL

☐ AChR antibody results

List Tried & Failed Therapies, including duration of treatment:

1)

2)

MEDICATION ORDERS

VYVGART™ (efgartigimod alfa-fcab)

☐ 10mg/kg IV over one hour weekly for 4 weeks* (Max Dose=1200mg) ☐ Other: _____

**Monitor patient for at least 1 hour after infusion.*

VYVGART HYTRULO™ (efgartigimod alfa and hyaluronidase-qvfc)

☐ 1008mg / 11,200 units SubQ once weekly for 4 weeks* ☐ Other: _____

**Monitor patient for at least 30 minutes after injection*

Subsequent treatment cycles to be at least 50 days from first dose of previous treatment

Refills: ☐ None ☐ X 6 months ☐ X 1 Year ☐ Other: _____

☐ RN to manage VAD per company protocol and administer ordered therapy per manufacturer guideline

☐ RN to access/start and deaccess/discontinue VAD as appropriate for therapy administration

☐ RN to flush and lock VAD/CVAD per company protocol

Other: _____

PREMEDICATION ORDERS

☐ Acetaminophen 650mg PO prior to infusion

☐ Other: _____

☐ Diphenhydramine 25mg PO prior to infusion

☐ Other: _____

EMERGENCY MEDICATIONS

☐ Administer the following medications as needed for infusion-related reactions per company protocol:

Adults (weight >40kg):

Diphenhydramine 25mg-50mg PO

Diphenhydramine 25mg-50mg slow IV push over 2-5 mins

Acetaminophen 325mg-650mg PO

Methylprednisolone 125mg slow IV push over at least 5 mins as tolerated

Epinephrine 0.3mg IM/SQ, may repeat x1

Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

Oxygen 1-6LPM continuous flow per nasal cannula or face mask, titrate to maintain SpO2 of 95-100% (A/C/AIS only)

Pediatrics (weight <40kg): (may adjust with weight changes)

Diphenhydramine 25mg PO

Diphenhydramine 25mg slow IV push over 2-5 mins

Acetaminophen 325mg PO

Methylprednisolone 40mg slow IV push over at least 5 mins as tolerated

Epinephrine 0.15mg (<30kg) or 0.3mg (>30kg) IM/SQ, may repeat x1

Sodium chloride 0.9% 500ml over 30-60 mins, may repeat x1 if hypotensive

PRESCRIBER INFORMATION

Prescriber Name: _____ NPI Number: _____

Office Phone: _____ Office Fax: _____

Prescriber Signature: _____ Date: _____