BIG SKY IV CARE
Infusion Pharmacy & Nursing
An Access Infusion Care Company

VUTISIRAN ORDERS (Amvuttra)

REFERRAL CHECKLIST
Please ensure the following information is included with your referral:
☐ Patient Demographics and Contact Information
☐ Patient Insurance Information
☐ Completed H&P and Progress Notes Including Medications and Allergies
☐ Recent Lab Results/Testing/Screenings Supporting the Diagnosis
☐ Completed and Signed Referral
☐ Cardiac Testing results (PYP scan, ECHO-if available)
Dx Codes: E85.4 Organ-limited amyloidosis E85.1 Neuropathic heredofamilial amyloidosis E85.82 Wild-type transthyretin-related (ATTR) amyloidosis
Including this information will minimize delays with prior authorizations and ensure timely treatment for your patient. Thank you!

©Big Sky IV Care
Print, complete, and fax completed form to (406) 752-0443

BIG SKY IV CARE
Infusion Pharmacy & Nursing
An Access Infusion Care Company

VUTISIRAN ORDERS (Amvuttra)

PATIENT INFORMATION	PRESCRIBER INFORMATION
Patient Name:	Prescriber Name:
Address:	NPI:
City, State, Zip:	Address:
Cell: Email:	City, State, Zip:
Ht: Wt: kg Last 4 of SSN: DOB:	Phone: Fax:
Allergies:	Contact Person:

REFERRAL STATUS ☐ New Referral ☐ Continuation Order ☐ Dose or Frequency Change

First dose? ☐ Yes ☐ No - Date of last injection:

DIAGNOSIS

Dx code and description:

REQUIRED DOCUMENTATION ☐ Signed order ☐ Demographics/insurance info ☐ H&P/labs/tests/progress notes supporting the Dx

☐ Monoclonal Protein levels	☐ Ensure patient is taking Vitamin A supplementation
☐ Cardiac PYP Scan	☐ Echocardiogram (if available)

Prior Therapies:

PRESCRIBING INFORMATION

LAB ORDERS

Frequency: Every Labs: ☐ CMP ☐ CBC w dif ☐ Other:

MEDICATION

Medication	Strength	Dose/Directions	Quantity
☒ **Vutisiran (Amvuttra)**	25mg/0.5mL	☒ 25 mg – Inject subcutaneously every 3 months	1 PFS Refills:

If not specified, order will be valid for 1 year from the date of signature.

OBSERVATION

Observe the patient for at least 30 minutes after the first infusion, no observation for subsequent infusions if no IRRs

NURSING INSTRUCTIONS/EMERGENCY MEDICATION

RN to manage VAD per company protocol and administer ordered therapy per manufacturer guidelines Post-Infusion:50ml0.9%NS to flush residual priming volume	PIV:Flush 3-10mL0.9%NS before/after PICC/Midline/Port:Flush 10mL0.9%NS before/after, then 3mLHeparin 10u/mL(PICC/Midline)/5mLHeparin 100u/mL(Port)final

Administer the following medications as needed for infusion-related reactions per company protocol:
Adults>40kg: Diphenhydramine25-50mgPO or IVP(2-5min); Acetaminophen325-650mgPO; Methylprednisolone125mgIVP(≥5min); Epinephrine0.3mgIM/SQ, may repeat×1; 0.9%NS500mLIV over 30-60 min, may repeat×1 for hypotension.
Pediatrics/Adults<40kg: Diphenhydramine25mgPO or IVP(2-5min); Acetaminophen325mgPO; Methylprednisolone40mgIVP(≥5min); Epinephrine0.15mg(<30kg)/0.3mg(>30kg)IM/SQ, may repeat×1; 0.9%NS500mLIV over 30-60min, may repeat×1 for hypotension.
Oxygen(AIC/AIS only):1-6L/min via NC/face mask; titrate to maintain $SpO_2$95-100%.

Prescriber Signature X: _____ **Date:** _____

☐ **Product Substitution Permitted** ☐ **Dispense as Written**

©Big Sky IV Care
Print, complete, and fax completed form to (406) 752-0443