BIG SKY IV CARE
Infusion Pharmacy & Nursing
An Access Infusion Care Company

C1 ESTERASE INHIBITOR ORDERS (Cinryze)

REFERRAL CHECKLIST
Please ensure the following information is included with your referral:
☐ Patient Demographics and Contact Information
☐ Patient Insurance Information
☐ Completed H&P and Progress Notes Including Medications and Allergies
☐ Recent Lab Results/Testing/Screenings Supporting the Diagnosis
☐ C1 Esterase Inhibitor Function Assay/C1-INH/C1q/C4/Genetic testing results
☐ Completed and Signed Referral
Including this information will minimize delays with prior authorizations and ensure timely treatment for your patient. Thank you!

©Big Sky IV Care

BIG SKY IV CARE
Infusion Pharmacy & Nursing
An Access Infusion Care Company

C1 ESTERASE INHIBITOR ORDERS (Cinryze)

PATIENT INFORMATION	PRESCRIBER INFORMATION
Patient Name:	Prescriber Name:
Address:	NPI:
City, State, Zip:	Address:
Cell: Email:	City, State, Zip:
Ht: Wt: kg Last 4 of SSN: DOB:	Phone: Fax:
Allergies:	Contact Person:

REFERRAL STATUS ☐ New Referral ☐ Continuation Order ☐ Dose or Frequency Change	
First dose? ☐ Yes ☐ No - Date of last infusion:	Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other:

DIAGNOSIS
☐ Dx Code/description:

REQUIRED DOCUMENTATION ☐ Signed order ☐ Demographics/insurance info ☐ H&P/labs/tests/progress notes supporting the Dx
Prior Therapies:
☐ C1 Esterase Inhibitor Function Assay/C1-INH/C1q/C4/Genetic testing results

PRESCRIBING INFORMATION

LAB ORDERS

Frequency: Every Labs: ☐ CMP ☐ CBCwdif ☐ Other:

MEDICATION

Medication	Strength	Dose/Directions	Quantity
☒ **C1 Esterase Inhibitors (Cinryze)**	500u/vial	☐ 1000units every 3-4 days	☐ 2 vial Refills:
		☐ units every	☐ # sufficient for dose Refills:
		Infuse IV at 1ml/min (dose should not exceed 80u/kg)	

If not specified, order will be valid for 1 year from the date of signature.

NURSING INSTRUCTIONS/EMERGENCY MEDICATION	
RN to manage VAD per company protocol and administer ordered therapy per manufacturer guidelines Post-Infusion:Use 50ml0.9%NS to flush residual priming volume	PIV:Flush 3-10mL0.9%NS before/after PICC/Midline/Port:Flush 10mL0.9%NS before/after, then 3mLHeparin 10u/mL(PICC/Midline)/ 5mLHeparin 100u/mL(Port) final

Administer the following medications as needed for infusion-related reactions per company protocol:
Adults>40kg: Diphenhydramine25-50mgPO or IVP(2-5min); Acetaminophen325-650mgPO; Methylprednisolone125mgIVP(≥5min); Epinephrine0.3mgIM/SQ, may repeat×1; 0.9%NS500mL IV over 30-60min, may repeat×1 for hypotension.
Pediatrics/Adults<40kg: Diphenhydramine25mgPO or IVP(2-5min); Acetaminophen325mgPO; Methylprednisolone40mgIVP(≥5min); Epinephrine0.15mg(<30kg)/0.3mg(>30kg)IM/SQ, may repeat×1; 0.9%NS500mL IV over 30-60min, may repeat×1 for hypotension.
Oxygen(AIC/AIS only):1-6L/min via NC/face mask; titrate to maintain SpO_2 95-100%.

Prescriber Signature X: _____ **Date:** _____

☐ **Product Substitution Permitted** ☐ **Dispense as Written**

©Big Sky IV Care
Print, complete, and fax completed form to (406) 752-0443