BIG SKY IV CARE
Infusion Pharmacy & Nursing
An Access Infusion Care Company

PEGLOTICASE ORDERS (Krystexxa)

REFERRAL CHECKLIST
Please ensure the following information is included with your referral:
☐ Patient Demographics and Contact Information
☐ Patient Insurance Information
☐ Completed H&P and Progress Notes Including Medications and Allergies
☐ Recent Lab Results/Testing/Screenings Supporting the Diagnosis
☐ Xray Reports Indicating Joint Damage from Gout
☐ Completed and Signed Referral
☐ Baseline Uric Acid Level, G6PD Screening
☐ Arrangements for Uric Acid Level lab Prior to Each Administration
☐ Completed and Signed Referral
Including this information will minimize delays with prior authorizations and ensure timely treatment for your patient. Thank you!

©Big Sky IV Care
Print, complete, and fax completed form to (406) 752-0443

BIG SKY IV CARE
Infusion Pharmacy & Nursing
An Access Infusion Care Company

PEGLOTICASE ORDERS (Krystexxa)

PATIENT INFORMATION	PRESCRIBER INFORMATION
Patient Name:	Prescriber Name:
Address:	NPI:
City, State, Zip:	Address:
Cell: Email:	City, State, Zip:
Height: Weight: kg Last 4 of SSN: DOB:	Phone: Fax:
Allergies:	Contact Person:

REFERRAL STATUS ☐ New Referral ☐ Continuation Order ☐ Dose or Frequency Change	
First dose? ☐ Yes ☐ No - Date of last infusion:	Line type: ☐ PIV ☐ PICC ☐ Port ☐ Other

DIAGNOSIS	
☐ M1A.00X0: Chronic gout with Tophus (Idiopathic)	Other Dx Code and description:

REQUIRED DOCUMENTATION	
☐ This signed order ☐ Patient demographics and insurance information ☐ H&P, labs, tests, progress notes supporting diagnosis	
• Prior Therapies:	☐ Negative G6PD screening – attach results
Methotrexate/folic acid supplementation start date: ☐ N/A-monotherapy	☐ Baseline Uric Acid Level (attach results):
☐ Patient has a standing Uric acid level lab order for 48 hours prior to each infusion. Will hold if >6mg/dL for two consecutive infusions.	

PRESCRIBING INFORMATION

LAB ORDERS

Frequency:Every Labs: ☐ CMP ☐ CBCwdif ☐ CRP ☐ Sed Rate ☐ ☐ Uric Acid level prior to each infusion

PREMEDICATIONS (administer 30 minutes prior to infusion, recommended by manufacturer)

Acetaminophen: ☐ 650mgPO ☐ 1000mgPO	Methylprednisolone: ☐ 100mgIV ☐ 125mgIV ☐ mgIV
Diphenhydramine: ☐ 25mgPO ☐ 50mgPO ☐ 25mgIV ☐ 50mgIV	Other: ☐

MEDICATION

Medication	Strength	Dose/Directions	Quantity
☒ **Pegloticase (Krystexxa)**	8mg/mL	☒ 8mg Infuse in 250ml 0.9%NS IV over 120 minutes every 2 weeks	1 vial Refills:

If not specified, order will be valid for 1 year from the date of signature.

OBSERVATION

Observe the patient for at least 60 minutes after the infusion.

NURSING INSTRUCTIONS/EMERGENCY MEDICATION

RN to manage VAD per company protocol and administer ordered therapy per manufacturer guidelines Post-Infusion:Use 50ml0.9%NS to flush residual priming volume	PIV:Flush 3-10mL0.9%NS before/after PICC/Midline/Port:Flush 10mL0.9%NS before/after, then 3mLHeparin 10u/mL(PICC/Midline)/ 5mLHeparin 100u/mL(Port) final

Administer the following medications as needed for infusion-related reactions per company protocol:
Adults>40kg: Diphenhydramine25-50mgPO or IVP(2-5min); Acetaminophen325-650mgPO; Methylprednisolone125mgIVP(≥5min); Epinephrine0.3mgIM/SQ, may repeat×1; 0.9%NS500mL IV over 30-60min, may repeat×1 for hypotension.
Pediatrics/Adults<40kg: Diphenhydramine25mgPO or IVP(2-5min); Acetaminophen325mgPO; Methylprednisolone40mgIVP(≥5min); Epinephrine0.15mg(<30kg)/0.3mg(>30kg)IM/SQ, may repeat×1; 0.9%NS500mL IV over 30-60min, may repeat×1 for hypotension.
Oxygen(AIC/AIS only):1-6L/min via NC/face mask; titrate to maintain SpO_2 95-100%.

Prescriber Signature X: _____ **Date:** _____

☐ **Product Substitution Permitted** ☐ **Dispense as Written**

©Big Sky IV Care
Print, complete, and fax completed form to (406) 752-0443