BIG SKY IV CARE
Infusion Pharmacy & Nursing
An Access Infusion Care Company

VELIGROTUG ORDERS (Lumvoa)

REFERRAL CHECKLIST
Please ensure the following information is included with your referral:
☐ Patient Demographics and Contact Information
☐ Patient Insurance Information
☐ Completed H&P and Progress Notes Including Medications and Allergies
☐ T3/T4/A1C/ documentation of failed high dose steroid
☐ Recent Lab Results/Testing/Screenings Supporting the Diagnosis
☐ Completed and Signed Referral
Including this information will minimize delays with prior authorizations and ensure timely treatment for your patient. Thank you!

©Big Sky IV Care
Print, complete, and fax completed form to (406) 752-0443

BIG SKY IV CARE
Infusion Pharmacy & Nursing
An Access Infusion Care Company

VELIGROTUG ORDERS (Lumvoa)

PATIENT INFORMATION	PRESCRIBER INFORMATION
Patient Name:	Prescriber Name:
Address:	NPI:
City, State, Zip:	Address:
Cell: Email:	City, State, Zip:
Ht: Wt: kg Last 4 of SSN: DOB:	Phone: Fax:
Allergies:	Contact Person:

REFERRAL STATUS ☐ New Referral ☐ Continuation Order ☐ Dose or Frequency Change	
First dose? ☐ Yes ☐ No - Date of last infusion:	Line type: ☐PIV ☐PICC ☐Port ☐Other

DIAGNOSIS	
☐ Dx Code:	

REQUIRED DOCUMENTATION ☐ Signed order ☐ Demographics and insurance info ☐ H&P, labs, tests, progress notes supporting dx	
☐Recent blood glucose or A1C results (attach results)	☐Confirmed Dx of TED from an Ophthalmologist (include documentation)
• Prior Therapies:	

PRESCRIBING INFORMATION

LAB ORDERS

Frequency: Every Labs: ☐CMP ☐CBCwdif ☐

PREMEDICATIONS

Acetaminophen: ☐650mgPO ☐1000mgPO	Methylprednisolone: ☐100mgIV ☐125mgIV ☐ mgIV
Diphenhydramine: ☐25mgPO ☐50mgPO ☐25mgIV ☐50mgIV	Other: ☐

MEDICATION

Medication	Strength	Dose/Directions	Quantity
☒**Veligrotug (Lumvoa)**	500mg/10 mL	☒**10mg/kg every 3 weeks x 5 doses** Infuse IV in 250ml 0.9%NS total volume over 45 minutes for the 1st infusion; then if tolerated, 30 minutes for all subsequent infusions	☒# sufficient for weight-based dose Refills: 4

Use current documented weight; round calculated dose to nearest vial size +/- 10% of dose.
If not specified, order will be valid for 1 year from the date of signature.

OBSERVATION

Observe the patient for at least 15 minutes after the infusion.

NURSING INSTRUCTIONS/EMERGENCY MEDICATION

RN to manage VAD per company protocol and administer ordered therapy per manufacturer guidelines Post-Infusion:Use 50ml0.9%NS to flush residual priming volume	PIV:Flush 3-10mL0.9%NS before/after PICC/Midline/Port:Flush 10mL0.9%NS before/after, then 3mLHeparin 10u/mL(PICC/Midline)/ 5mLHeparin 100u/mL(Port) final

Administer the following medications as needed for infusion-related reactions per company protocol:
Adults>40kg: Diphenhydramine25-50mgPO or IVP(2-5min); Acetaminophen325-650mgPO; Methylprednisolone125mgIVP(≥5min); Epinephrine0.3mgIM/SQ, may repeat×1; 0.9%NS500mL IV over 30-60min, may repeat×1 for hypotension.
Pediatrics/Adults<40kg: Diphenhydramine25mgPO or IVP(2-5min); Acetaminophen325mgPO; Methylprednisolone40mgIVP(≥5min); Epinephrine0.15mg(<30kg)/0.3mg(>30kg)IM/SQ, may repeat×1; 0.9%NS500mL IV over 30-60min, may repeat×1 for hypotension.
Oxygen(AIC/AIS only):1-6L/min via NC/face mask; titrate to maintain SpO_2 95-100%.

Prescriber Signature X: _____ **Date:** _____

☐**Product Substitution Permitted** ☐**Dispense as Written**

©Big Sky IV Care
Print, complete, and fax completed form to (406) 752-0443